arXiv:1411.5032v1 [hep-ph] 18 Nov 2014

R symmetries and a heterotic MSSM

Rolf Kappl, Hans Peter Nilles, Matthias Schmitz

Bethe Center for Theoretical Physics
and
Physikalisches Institut der Universität Bonn
Nussallee 12, 53115 Bonn, Germany

Abstract

We employ powerful techniques based on Hilbert- and Gröbner bases to analyze particle physics models derived from string theory. Individual models are shown to have a huge landscape of vacua that differ in their phenomenological properties. We explore the (discrete) symmetries of these vacua, the new R symmetry selection rules and their consequences for moduli stabilization.

1 Introduction

Recently new results on R symmetries in heterotic orbifolds have been obtained [1, 2]. We discuss the implications of these results on phenomenologically appealing models like the ones obtained in [3–5]. Our purpose is twofold. On the one hand we show that with the new R symmetries successful models remain. On the other hand we employ techniques developed in [6] to study the vacuum configuration of a given model. These techniques enable us to determine the structure of the superpotential in form of building monomials. The advantage to previous attempts is, that one can immediately see which standard model singlets induce which couplings. We further show that this approach is also useful for general particle physics models with continuous or discrete symmetries. All of these symmetries lead to Diophantine equations whose solutions are given by so-called Hilbert basis elements. After the determination of the superpotential we search for suitable SUSY preserving minima of our model. We find that some of the standard model singlets will get stabilized at a non-trivial value. In contrast to [7] many singlets will have flat directions. This can be understood in terms of remnant symmetries.

The paper is outlined as follows. In section 2 we show how symmetries result in Diophantine equations and how these can be solved to determine the structure of the underlying physical model. In section 3 we briefly review new results for R symmetries in the heterotic orbifold context obtained in [1, 2] and comment on their consequences. We finally apply the developed techniques to a phenomenologically appealing model in section 4 and summarize our results.

2 Allowed couplings and Hilbert bases

It has been outlined in [6] how to use Hilbert bases to compute a basis of all allowed monomials in the superpotential. In this section we will use this technique to derive which couplings are allowed by the R symmetries and the other well known string selection rules. We will explain the approach by an example from flavor model building [8] and apply the technique to our concrete string model.

2.1 Symmetries and Diophantine equations

Given some continuous or discrete symmetries one is usually interested in the question which couplings are allowed by the given symmetries. Let us assume a theory with a U(1) symmetry and fields ϕ_i with charges q_i under the symmetry. An example for an allowed monomial would be

$$W \supset \phi_1\phi_2\phi_3^2 \qquad \Leftrightarrow \qquad q_1 + q_2 + 2q_3 = 0\,. \tag{1}$$

We can generalize this to

$$W \supset \phi_1^{n_1} \dots \phi_M^{n_M} \qquad \Leftrightarrow \qquad q^T \cdot n = 0\,, \tag{2}$$

where $q^T = (q_1, \dots, q_M)$ and $n^T = (n_1, \dots, n_M) \in \mathbb{N}_0^M$. Because of the restriction $n_i \in \mathbb{N}_0$ this is called a Diophantine equation. That means every Abelian gauge symmetry leads

naturally to a Diophantine equation. It is not possible to give an analytical basis for all solution vectors n because the natural numbers only form a monoid. Nevertheless it is possible to algorithmically determine a basis. We can write

$$n = \sum_i \alpha_i x_i, \qquad x_i \in \mathcal{H}, \qquad \alpha_i \in \mathbb{N}_0\,. \tag{3}$$

Note that the total number of Hilbert basis elements x_i cannot be determined analytically. The basis spanned by the elements x_i is called Hilbert basis and several freely available programs [9, 10] exist to determine it. This was shown and outlined in more detail in [6]. Let us consider an example from flavor model building, namely the model discussed in [8]. Some of its fields and their respective charges under a large number of discrete symmetries are given in table 1. We are interested in Yukawa couplings involving two fields $\mathcal{T}_i$ and a

	$\mathbb{Z}_2$	$\mathbb{Z}_4$	$\mathbb{Z}_4$	$\mathbb{Z}_4$	$\mathbb{Z}_7$	$\mathbb{Z}_7$	$\mathbb{Z}_9$	$\mathbb{Z}_2$
H_5	0	0	0	0	0	0	0	0
$\mathcal{T}_1$	0	0	3	0	6	0	0	1
$\mathcal{T}_2$	0	0	0	0	6	0	0	1
$\mathcal{T}_3$	0	0	0	0	0	0	0	1
θ_1	0	0	0	0	1	0	0	0
θ_2	0	0	1	0	0	0	0	0
θ_3	0	2	0	0	0	6	5	0
θ_4	0	0	0	0	0	0	5	0

Table 1: Charges of the fields considered in the flavor model discussed in [8].

neutral Higgs field H_5. We can compute the Hilbert basis $\mathcal{H}$ for these couplings with the help of `normaliz` [9, 11] and find the following elements in this model

$$\begin{aligned} W = H_5 \big(&\mathcal{T}_1\mathcal{T}_1(\theta_1^2\theta_2^2 + \theta_1^2\theta_2^6 + \theta_1^9\theta_2^2 + \theta_1^2\theta_2^2\theta_4^9 + \ldots) + \mathcal{T}_1\mathcal{T}_2(\theta_1^2\theta_2 + \ldots) + \mathcal{T}_2\mathcal{T}_2(\theta_1^2 + \ldots) \\ &+\mathcal{T}_2\mathcal{T}_3(\theta_1 + \ldots) + \mathcal{T}_3\mathcal{T}_3(1 + \ldots)\big)\,. \end{aligned} \tag{4}$$

This result is in agreement with the superpotential given in [8] and shows the applications of our approach. The dots denote monomials which consist of higher order couplings in the flavon fields θ_i. Note that while we considered only Abelian symmetries here, it would also be interesting to consider Hilbert bases for non-Abelian discrete symmetries like the ones discussed for example in [12].

2.2 Diophantine equations and R symmetries

R symmetries result in inhomogeneous Diophantine equations. In this section we will already have a concrete string model in mind (benchmark model 1 from [4]). Nonetheless, the results are applicable to any model with R symmetries. Our approach also deviates slightly from [6]. Let us assume that our theory gives rise to the R symmetry[1] $\mathbb{Z}_6^R \times \mathbb{Z}_3^R \times \mathbb{Z}_2^R$. This is for example the case for models in the mini-landscape and has

[1] As outlined, for example, in [7] the charges under these symmetries are not integers, which means that a redefinition seems to be natural. Nevertheless we keep this form for easier comparison with the literature.

been reconsidered recently in [1,2] which is one motivation of our work. Let us denote the charges under these symmetries by $R^T = (R^1, R^2, R^3)$. Thus the R symmetry of the model will result in the constraint

$$\sum_i R_i^T = (-1,-1,-1) \mod (6,3,2)\,, \tag{5}$$

where R_i denotes the R charge of a field ϕ_i and we assume that the superpotential transforms with R charge one. Let us illustrate our approach with an example. If we want to know which singlets $\tilde{s}_i$ induce the Yukawa term

$$W \supset \bar{\phi}_1 q_1 \bar{u}_1 f(\tilde{s}_i)\,, \tag{6}$$

we have to solve the inhomogeneous system of Diophantine equations

$$\hat{R}\cdot n = \begin{pmatrix}-1\\-1\\-1\end{pmatrix} \mod \begin{pmatrix}6\\3\\2\end{pmatrix}\,, \qquad n_i \in \mathbb{N}_0 \quad \forall i\,, \tag{7}$$

with

$$\hat{R} = \begin{pmatrix}R_1 & \dots & R_{M+3}\end{pmatrix}\,, \qquad R_i^T = (R_i^1, R_i^2, R_i^3)\,, \tag{8}$$

where M is the number of different standard model singlets in our model. During this discussion we use the notation from [4] and call the up type Higgs $\bar{\phi}_1$ and the down type Higgs ϕ_1. Further, as in section 2.1, n_i denotes the exponent of a given field

$$W \supset \bar{\phi}_1^{n_1} q_1^{n_2} \bar{u}_1^{n_3} (\tilde{s}_1)^{n_4} (\tilde{s}_2)^{n_5} \dots (\tilde{s}_M)^{n_{M+3}}\,. \tag{9}$$

Only solutions with $n_1 = n_2 = n_3 = 1$ are physical, therefore we can simplify this equation system by substituting

$$\tilde{R} = R_1 + R_2 + R_3 + (1,1,1)^T\,. \tag{10}$$

This results in a homogeneous system of equations with less variables

$$\begin{pmatrix}\tilde{R} & R_4 & \dots & R_{M+3}\end{pmatrix}\cdot\begin{pmatrix}n_1\\n_4\\\vdots\\n_{M+3}\end{pmatrix} = \begin{pmatrix}0\\0\\0\end{pmatrix} \mod \begin{pmatrix}6\\3\\2\end{pmatrix}\,, \qquad n_i \in \mathbb{N}_0 \quad \forall i\,. \tag{11}$$

We have discussed how to find solutions to such homogeneous Diophantine equation systems in section 2.1. After determining a Hilbert basis $\mathcal{H}$ one can further split it to obtain physically viable solutions. Elements with $n_1 = 1$ are assigned to $\mathcal{H}_{\text{inhom}} \subset \mathcal{H}$ and basis elements with $n_1 = 0$ are assigned to $\mathcal{H}_{\text{hom}} \subset \mathcal{H}$. All physical solutions to equation (11) are then given by the vectors

$$x = x_{\text{inhom}}(1 + x_{\text{hom}} + x_{\text{hom}}^2 + \dots)\,, \tag{12}$$

where $\dots$ denote higher powers in x_{hom}. One has to take all possible combinations of elements $x_{\text{inhom}} \in \mathcal{H}_{\text{inhom}}$ and $x_{\text{hom}} \in \mathcal{H}_{\text{hom}}$. For practical purposes it seems reasonable to truncate the solution at some finite order in the fields ϕ_i.

2.3 The superpotential to all orders

We can now determine all monomials which give W for the string model under consideration. We follow the approach of [6] and view all string selection rules as gauge and discrete $\mathbb{Z}_N$ symmetries. Note that the space group is an infinite discrete non-Abelian group. However for $\mathbb{Z}_6$-II models, it is possible to rephrase it in terms of the finite discrete group[2] $\mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_2'$ [13].

Our model enjoys the symmetries $\mathbb{Z}_6^R \times \mathbb{Z}_3^R \times \mathbb{Z}_2^R \times \mathbb{Z}_6 \times \mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_2' \times \mathrm{U}(1)^8$. The R symmetries are remnants of the internal Lorentz symmetries, the $\mathbb{Z}_6$ symmetry results from the so-called point group selection rule, whereas the space group selection rule results in $\mathbb{Z}_3 \times \mathbb{Z}_2 \times \mathbb{Z}_2'$. In addition we have to take care of the gauge symmetries resulting from the $\mathrm{E}_8 \times \mathrm{E}_8$. The non-Abelian symmetries can be discussed along the lines of [6] and we focus on the Cartan generators leading to the U(1) factors given here. With the discussion of the R symmetries from section 2.2 we get the Diophantine equation system

$$\begin{pmatrix} \tilde{R}^1 & R_4^1 & \dots & R_{M+3}^1 \\ \tilde{R}^2 & R_4^2 & \dots & R_{M+3}^2 \\ \tilde{R}^3 & R_4^3 & \dots & R_{M+3}^3 \\ q_1^{\mathbb{Z}_6} + q_2^{\mathbb{Z}_6} + q_3^{\mathbb{Z}_6} & q_4^{\mathbb{Z}_6} & \dots & q_{M+3}^{\mathbb{Z}_6} \\ q_1^{\mathbb{Z}_3} + q_2^{\mathbb{Z}_3} + q_3^{\mathbb{Z}_3} & q_4^{\mathbb{Z}_3} & \dots & q_{M+3}^{\mathbb{Z}_3} \\ q_1^{\mathbb{Z}_2} + q_2^{\mathbb{Z}_2} + q_3^{\mathbb{Z}_2} & q_4^{\mathbb{Z}_2} & \dots & q_{M+3}^{\mathbb{Z}_2} \\ q_1^{\mathbb{Z}_2'} + q_2^{\mathbb{Z}_2'} + q_3^{\mathbb{Z}_2'} & q_4^{\mathbb{Z}_2'} & \dots & q_{M+3}^{\mathbb{Z}_2'} \\ q_1^1 + q_2^1 + q_3^1 & q_4^1 & \dots & q_{M+3}^1 \\ \vdots & & & \vdots \\ q_1^8 + q_2^8 + q_3^8 & q_4^8 & \dots & q_{M+3}^8 \end{pmatrix} \cdot \begin{pmatrix} n_1 \\ n_4 \\ \vdots \\ n_{M+3} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \mod \begin{pmatrix} 6 \\ 3 \\ 2 \\ 6 \\ 3 \\ 2 \\ 2 \\ 0 \\ \vdots \\ 0 \end{pmatrix} . \tag{13}$$

We use `normaliz` [9, 11] to compute the Hilbert basis $\mathcal{H}$ of this Diophantine equation system. Compared to other systems like `4ti2` [10] it has the advantage that the algorithm used is suited better for our problem [14] and is also able to deal with congruences. We use the so-called "primal" algorithm [14] which is much faster in our case then the so-called "dual" algorithm which is implemented in `4ti2`.

To make our results comparable with the results from the literature we reconsider benchmark model 1 from [4] in the following. We apply the rederived R symmetries which have been determined in [1, 2] and which we review in section 3. We consider a vacuum configuration in which we give the 14 standard model singlets

$$\tilde{s} = \{h_1, h_2, h_3, h_4, s_3^0, s_4^0, s_9^0, s_{10}^0, s_{12}^0, s_{21}^0, s_{24}^0, s_{28}^0, s_{29}^0, s_{30}^0\} \tag{14}$$

a vacuum expectation value (VEV). To get the superpotential W of these fields to all orders, we can consider which fields induce the μ-term. This is possible because $\phi_1\bar{\phi}_1$ forms a complete singlet under all symmetries in this model [4, 15, 16]. We find that up to order 10 in standard model singlets $\tilde{s}_i$ the superpotential is given by

$$\begin{aligned} W &= (M_{1,\mathrm{inhom}} + M_{2,\mathrm{inhom}} + M_{3,\mathrm{inhom}})(1 + M_{1,\mathrm{hom}} + M_{2,\mathrm{hom}}) \\ &= ((s_3^0 s_4^0 + s_9^0 s_{10}^0) s_{29}^0 + h_3 h_4 s_{21}^0 s_{31}^0)(1 + h_1 h_2 s_{21}^0 s_{24}^0 s_{28}^0 s_{31}^0 + (s_{21}^0 s_{31}^0)^3)\,, \end{aligned} \tag{15}$$

[2] There is an additional rule for some exceptional cases in the G_2 orbifold plane [13]. We have not been able to write it as a discrete symmetry and checked the invariance of the basis elements separately case by case a posterior.

with

$$\begin{aligned} M_{1,\text{inhom}} &= s_3^0 s_4^0 s_{29}^0\,, & M_{1,\text{hom}} &= (s_{21}^0 s_{31}^0)^3\,, \\ M_{2,\text{inhom}} &= s_9^0 s_{10}^0 s_{29}^0, & M_{2,\text{hom}} &= h_1 h_2 s_{21}^0 s_{24}^0 s_{28}^0 s_{31}^0\,, \\ M_{3,\text{inhom}} &= h_3 h_4 s_{21}^0 s_{31}^0\,. \end{aligned} \tag{16}$$

The monomials $M_{i,\text{inhom}}$ are given by their exponent vectors $x_{i,\text{inhom}}$ which are elements of the corresponding part of the Hilbert basis $x_{i,\text{inhom}} \in \mathcal{H}_{\text{inhom}}$. The same is true for the homogeneous part. In principle we know all Hilbert basis elements and therefore all monomials which means we know the exact form of W to all orders. We also see the manifestation of the D_4 symmetry [17,18] present in this model which relates $M_{1,\text{inhom}}$ and $M_{2,\text{inhom}}$. There are two doublets

$$\tilde{D}_1 = (s_3^0, s_9^0)\,, \qquad \tilde{D}_2 = (s_4^0, s_{10}^0)\,, \tag{17}$$

which result in two invariants

$$\tilde{D}_1 \cdot \tilde{D}_2^T s_{29}^0 = (s_3^0 s_4^0 + s_9^0 s_{10}^0) s_{29}^0 = M_{1,\text{inhom}} + M_{2,\text{inhom}}\,. \tag{18}$$

3 *R* symmetries in heterotic orbifolds

In this section we briefly review the origin of discrete R symmetries in heterotic orbifolds [1,2,19]. These symmetries arise from automorphisms of the orbifold space group. For the case of the $\mathbb{Z}_6$-II orbifold the generators R^α can be written in terms of the Cartan generators of the internal SO(6) by means of twist vectors v_α according to $R^\alpha = \mathrm{e}^{2\pi\mathrm{i}(v_\alpha^1 J_{45} + v_\alpha^2 J_{67} + v_\alpha^3 J_{78})}$. They are given by

$$v_1 = \left(\tfrac{1}{6},\, 0,\, 0\right)\,, \quad v_2 = \left(0,\, \tfrac{1}{3},\, 0\right)\,, \quad v_3 = \left(0,\, 0,\, \tfrac{1}{2}\right)\,. \tag{19}$$

In order to determine the charges of the corresponding fields with respect to the R symmetries, the transformation behavior of the string states under these generators needs to be worked out.

String states of heterotic orbifolds are characterized by their left- and right-moving momenta p_{sh} and q_{sh}, their left-moving oscillator excitations $\tilde{\alpha}$ and their locus. As the properties of the strings, described by these quantum numbers, are mainly independent of each other it is customary to write such states as

$$|\Psi\rangle = \tilde{\alpha}^{\mathcal{N}_\text{L}}\, (\tilde{\alpha}^*)^{\bar{\mathcal{N}}_\text{L}}\; |p_{\text{sh}}\rangle \otimes |q_{\text{sh}}\rangle \otimes |\text{locus}\rangle\,. \tag{20}$$

Here $\mathcal{N}_\text{L}$ and $\bar{\mathcal{N}}_\text{L}$ count the number of holomorphic and anti-holomorphic oscillators of the state. Recall that twisted strings have their center of mass momentum localized at fixed points of the space group S. The constructing element $g \in S$ of a fixed point z_f is defined by $gz_f = z_f$. Untwisted strings can propagate freely and hence the factor describing the localization in (20) is absent for such states.

Note that we know the action of the generators (19) on the covering space $\mathbb{C}^3$ of the orbifold only. However, on $\mathbb{C}^3$, there are infinitely many copies of each fixed point, which are related to z_f by the action of space group elements h that do not commute with g. The corresponding constructing elements form the conjugacy class $[g] = \{hgh^{-1} | h \in S\}$

of g. Therefore, if we denote the locus of a string state located at the fixed point z_f with constructing element g by $|g\rangle$, a physical state corresponds to the linear combination

$$|\text{locus}\rangle = \sum_{g' \in [g]} \mathrm{e}^{2\pi \mathrm{i} \tilde{\gamma}(g')} \, |\Psi_{g'}\rangle \ . \tag{21}$$

The phases $\tilde{\gamma}$ are fixed by the requirement of invariance of the string states under elements h that do not commute with g, such that

$$\gamma(g,h) = \tilde{\gamma}(g) - \tilde{\gamma}(hgh^{-1}) = -p_{\text{sh}} \cdot V_h + v_h \cdot (q_{\text{sh}} - \mathcal{N}_{\text{L}} + \bar{\mathcal{N}}_{\text{L}}) \mod 1 \, . \tag{22}$$

Here v_h denotes the twist vector of the space group element h and V_h denotes its gauge embedding.

The crucial observation is now, that the rotations (19) map each space group element g to a conjugate one, i.e. for each g there exists an h_g such that $R(g) = h_g \, g \, h_g^{-1}$. Therefore,the transformation behavior of the linear combination (21) is given by

$$|\text{locus}\rangle \overset{R}{\longmapsto} \mathrm{e}^{-2\pi \mathrm{i} \gamma(g,h_g)} \, |\text{locus}\rangle \ . \tag{23}$$

For the other parts of the string states the transformation behavior follows from the transformation of the space time coordinates. A string state (20) transforms according to

$$|\Psi\rangle \overset{R}{\longmapsto} \exp\left[2\pi \mathrm{i} \, v \cdot \left(q_{\text{sh}} - \mathcal{N}_{\text{L}} + \bar{\mathcal{N}}_{\text{L}}\right) - 2\pi \mathrm{i} \, \gamma(g,h_g)\right] |\Psi\rangle \tag{24}$$

under an R symmetry generator R with shift vector v. For the case we are considering here, asking string correlators corresponding to superpotential terms to transform trivially under the R symmetries one arrives at the R charge selection rules [1, 2]

$$\begin{aligned}
\sum_{i=1}^{L} R_i^1 &= \sum_{i=1}^{L} \left[q_{\text{sh}\,i}^{(\text{boson})\,1} - \mathcal{N}_{\text{L}\,i}^1 + \bar{\mathcal{N}}_{\text{L}\,i}^1 - 6\gamma(g_i,\, h_{g_i}) \right] = -1 \mod 6 \, , \\
\sum_{i=1}^{L} R_i^2 &= \sum_{i=1}^{L} \left[q_{\text{sh}\,i}^{(\text{boson})\,2} - \mathcal{N}_{\text{L}\,i}^2 + \bar{\mathcal{N}}_{\text{L}\,i}^2 - 3\gamma(g_i,\, h_{g_i}) \right] = -1 \mod 3 \, , \\
\sum_{i=1}^{L} R_i^3 &= \sum_{i=1}^{L} \left[q_{\text{sh}\,i}^{(\text{boson})\,3} - \mathcal{N}_{\text{L}\,i}^3 - \bar{\mathcal{N}}_{\text{L}\,i}^3 - 2\gamma(g_i,\, h_{g_i}) \right] = -1 \mod 2 \, .
\end{aligned} \tag{25}$$

Here we have made use of the fact that the internal right-moving momenta q_{sh} of bosons and fermions within the same chiral multiplet are related by a shift of $\frac{1}{2}$. We denote the momentum of the respective bosons by $q_{\text{sh}}^{(\text{boson})}$ and define the R charge of a string state to be that of the bosonic component of the chiral multiplet.

We modified the program `orbifolder` [20] to incorporate the outlined R symmetries. The result for a concrete model can be found in appendix A.

3.1 Differences to previous results

Differences due to the new R symmetries arise already at order 3 in the superpotential. Several new terms appear which were forbidden by the old rules. In the following we show

	$R^{1,\text{old}}$	$R^{2,\text{old}}$	$R^{3,\text{old}}$	$R^{1,\text{new}}$	$R^{2,\text{new}}$	$R^{3,\text{new}}$
s^0_{13}	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$
s^0_{14}	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$
s^0_{30}	$-\frac{5}{3}$	$-\frac{1}{3}$	0	$\frac{4}{3}$	$-\frac{1}{3}$	0

Table 2: Differences between the R charges for some singlet fields.

a simple example. We look at the allowed couplings in the so-called benchmark model 1A of [4]. The R charges of the considered fields are displayed in table 2. We immediately see that the coupling $s^0_{13}s^0_{14}s^0_{30}$ was forbidden by the old rules but is allowed under the new R symmetries.

This has important phenomenological implications. If all fields s^0_{13}, s^0_{14} and s^0_{30} get a VEV in a chosen configuration, the superpotential W also gets a VEV. As already discussed, the superpotential in such models is linked to the μ-term. Therefore, in this configuration, the μ-term will be generically too large. We will discuss this issue for our VEV configuration later in more detail. There are also couplings forbidden by the new R symmetries which were allowed by the old symmetries. The first example occurs at order 4 in the superpotential.

4 Phenomenological properties

In this section we want to briefly comment on the phenomenological properties of the considered model. Instead of doing a complete scan over different vacua we stick to benchmark model 1 (the field content of this model can be found in appendix A). A full scan over different configurations is beyond the scope of this work. Using the vacuum configuration given in equation (14) we obtain the following features.

4.1 *F*-flatness, Gröbner bases and *D*-flatness

To analyze F-flatness we use techniques and methods known in computational algebraic geometry and discussed in high energy physics for example in [21] and in string theory in [22, 23]. We are looking for solutions to the equation system

$$F_i = \frac{\partial W}{\partial \tilde{s}_i} = 0, \qquad \forall i\,. \tag{26}$$

To find a solution we truncate the superpotential W at a given order. We take W up to order 10 in standard model singlets which was given in equation (15) and set all coupling coefficients to unity for simplicity. In principle these coefficients are calculable functions of the geometric moduli of the compactification. This dependence can be used to stabilize the geometric moduli [16].

We use `Singular` [24] to compute the Gröbner basis of the ideal generated by the F-term equations. Afterwards we compute the primary decomposition and search for F-flat solutions $F_i = 0$. We find only one non-trivial branch of solutions. Trivial solutions

$\langle \tilde{s}_i \rangle = 0$, would violate our assumption that all fields $\tilde{s}_i$ given in equation (14) obtain a non-vanishing VEV. The only non-trivial branch can be solved, for example, by

$$\left\langle s_{28}^0 \right\rangle = -\frac{1 + \left\langle s_{21}^0 \right\rangle^3 \left\langle s_{31}^0 \right\rangle^3}{\langle h_1 \rangle \langle h_2 \rangle \left\langle s_{21}^0 \right\rangle \left\langle s_{24}^0 \right\rangle \left\langle s_{31}^0 \right\rangle}\,, \qquad \left\langle s_{29}^0 \right\rangle = -\frac{\langle h_3 \rangle \langle h_4 \rangle \left\langle s_{21}^0 \right\rangle \left\langle s_{31}^0 \right\rangle}{\left\langle s_3^0 \right\rangle \left\langle s_4^0 \right\rangle + \left\langle s_9^0 \right\rangle \left\langle s_{10}^0 \right\rangle}\,, \tag{27}$$

which results in $F_i = \langle W \rangle = 0$. That means we can solve all 14 F-term equations simultaneously by fixing only two VEVs. This is nearly the opposite behavior to the one discussed in [7] where a remnant $\mathbb{Z}_4^R$ symmetry [25] has been used to restrict the superpotential[3]. There it seems to be more fertile to look for minima in which all singlets get fixed by the F-term equations. It is interesting to study how the symmetries of the superpotential determine the solution structure of the F-term equations. In our case many F-term equations are degenerate because of the remnant D_4 symmetry. The necessary breaking of this symmetry at a lower scale (see section 4.2) can therefore be potentially used to stabilize additional moduli. The detailed study of this mechanism is beyond the scope of this work. At this stage we are satisfied with finding a consistent, non-trivial solution.

We also would like to emphasize that the μ-term in the minima determined above, vanishes because $\langle W \rangle = 0$. This is well known to be related to approximate R symmetries [15]. As we know the superpotential not only to order 10 in singlets, but to all orders, it seems to be possible to address the question of F-flatness to all orders. It is effective to attack this problem in terms of the Hilbert basis monomials. In this way, the given non-trivial solution can be extended to all orders. We can split the superpotential according to the Hilbert basis monomials into two pieces

$$W(\tilde{s}_i) = W_{\mathrm{inhom}}(\tilde{s}_i)(1 + W_{\mathrm{hom}}(\tilde{s}_i))\,. \tag{28}$$

Here W_{inhom} is given by the linear combination of all inhomogeneous Hilbert basis monomials $M_{i,\mathrm{inhom}}$. Furthermore W_{hom} denotes all possible combinations of homogeneous monomials $M_{i,\mathrm{hom}}$. The F-term equations in this picture are given by

$$F_i = \frac{\partial W_{\mathrm{inhom}}}{\partial \tilde{s}_i}(1 + W_{\mathrm{hom}}) + W_{\mathrm{inhom}} \frac{\partial W_{\mathrm{hom}}}{\partial \tilde{s}_i}\,, \qquad \forall i\,. \tag{29}$$

It is obvious that all F-term equations vanish if $W_{\mathrm{inhom}}(\tilde{s}_i) = 0$ and $1 + W_{\mathrm{hom}}(\tilde{s}_i) = 0$ simultaneously. Our concrete solution at order 10 is exactly of this kind. This behavior is not limited to any order and in general such a solution will exist for the superpotential at higher order. The disadvantage that generically only two fields are fixed in this solution branch nevertheless remains. Different strategies to find minima like the ones considered in [28, 29] may also help to address the computational difficulties.

Let us also comment on D-flatness. It is possible to satisfy D-flatness along the lines of [30, 31]. We have explicitly checked that we can cancel the Fayet-Iliopoulos term by a monomial which carries negative charge under the anomalous U(1). D-flatness will also help to fix some VEVs (see for example [7]) but in our concrete example some flat directions remain.

[3]Further applications of this symmetry to flavor model building are discussed in [26], whereas the relation to R parity violation is studied in [27].

4.2 Yukawa couplings

For the Yukawa interactions we obtain

$$W_{\text{Yuk}} = Y_u(\tilde{s}_i) q \bar{u} \bar{\phi}_1 + Y_d(\tilde{s}_i) q \bar{d} \phi_1 + Y_e(\tilde{s}_i) l \bar{e} \phi_1 \,, \tag{30}$$

where the Yukawa matrices $Y_i(\tilde{s}_i)$ depend on the singlet fields $\tilde{s}_i$ to which we have assigned VEVs (see equation (14)). We computed for each coupling the corresponding Hilbert basis and found that, at lowest order in singlets, the structure is

$$Y_u = \begin{pmatrix} M_1 & M_2 & M_3 + M_4 \\ M_2 & M_1 & M_5 + M_6 \\ M_7 + M_8 + M_9 + M_{10} & M_{11} + M_{12} + M_{13} + M_{14} & 1 \end{pmatrix} , \tag{31}$$

with

$$\begin{aligned} M_1 &= h_1 h_3 s_4^0 s_{21}^0 s_{29}^0 s_{31}^0 \,, & M_2 &= h_1 h_3 s_{10}^0 s_{21}^0 s_{29}^0 s_{31}^0 \,, \\ M_3 &= s_3^0 M_2 \,, & M_4 &= s_9^0 M_1 \,, \\ M_5 &= s_3^0 M_1 \,, & M_6 &= s_9^0 M_2 \,, \\ M_7 &= s_{12}^0 M_1 \,, & M_8 &= s_3^0 s_4^0 s_{10}^0 s_{21}^0 (s_{29}^0)^2 s_{31}^0 \,, \\ M_9 &= s_9^0 (s_{10}^0)^2 s_{21}^0 (s_{29}^0)^2 s_{31}^0 \,, & M_{10} &= (s_4^0)^2 s_9^0 s_{21}^0 (s_{29}^0)^2 s_{31}^0 \,, \\ M_{11} &= s_{12}^0 M_2 \,, & M_{12} &= s_4^0 s_9^0 s_{10}^0 s_{21}^0 (s_{29}^0)^2 s_{31}^0 \,, \\ M_{13} &= s_3^0 (s_{10}^0)^2 s_{21}^0 (s_{29}^0)^2 s_{31}^0 \,, & M_{14} &= s_3^0 (s_4^0)^2 s_{21}^0 (s_{29}^0)^2 s_{31}^0 \,. \end{aligned} \tag{32}$$

Further, for the down quarks and leptons

$$Y_d = \begin{pmatrix} M_1 & M_2 & M_3 + M_4 \\ M_2 & M_1 & M_5 \\ M_6 & M_7 & M_8 + M_9 \end{pmatrix} , \qquad Y_e = \begin{pmatrix} M_1 & M_2 & M_{10} + M_{11} \\ M_2 & M_1 & M_{12} \\ M_{13} & M_{14} & M_{15} + M_{16} \end{pmatrix} , \tag{33}$$

with

$$\begin{aligned} M_1 &= h_1 h_2 s_9^0 s_{12}^0 s_{29}^0 \,, & M_2 &= h_1 h_2 s_3^0 s_{12}^0 s_{29}^0 \,, \\ M_3 &= s_9^0 M_1 \,, & M_4 &= s_3^0 M_2 \,, \\ M_5 &= s_3^0 M_1 \,, & M_6 &= s_{12}^0 s_{21}^0 M_1 \,, \\ M_7 &= s_{12}^0 s_{21}^0 M_2 \,, & M_8 &= s_9^0 s_{12}^0 s_{21}^0 M_1 \,, \\ M_9 &= s_3^0 s_{12}^0 s_{21}^0 M_2 \,, & M_{10} &= (s_9^0)^2 s_{12}^0 s_{21}^0 s_{29}^0 \,, \\ M_{11} &= (s_3^0)^2 s_{12}^0 s_{21}^0 s_{29}^0 \,, & M_{12} &= s_3^0 s_4^0 s_{12}^0 s_{21}^0 s_{29}^0 \,, \\ M_{13} &= s_{12}^0 M_1 \,, & M_{14} &= s_{12}^0 M_2 \,, \\ M_{15} &= s_{12}^0 M_{10} \,, & M_{16} &= s_{12}^0 M_{11} \,. \end{aligned} \tag{34}$$

Let us note that, as expected, the top quark coupling is of order one. The reason for this behavior is the connection of the coupling to the higher dimensional gauge coupling [32]. Thus a realistic top quark mass in this model is guaranteed.

As a consequence of the localization of the first two generations in the extra dimensional space, these fields form a doublet under the D_4 symmetry [17, 18]. This manifests itself in the appearance of the monomials M_1 and M_2 in the Yukawa matrices. However, this symmetry needs to be broken at a lower scale to explain the different masses between the first and second generation [33].

4.3 Additional features

Also with the new R symmetries all exotics can be made massive. We will discuss one example in detail and skip the details for the other exotics. As can be seen from table 3 the R^3 charges of y_1 and y_2 have changed under the new R symmetry. Nevertheless the

	$R^{1,\text{old}}$	$R^{2,\text{old}}$	$R^{3,\text{old}}$	$R^{1,\text{new}}$	$R^{2,\text{new}}$	$R^{3,\text{new}}$
y_1	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$
y_2	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$

Table 3: Differences between the R charges of y_1 and y_2.

mass matrix does not change, because both fields always appear in pairs and we have

$$\sum_i R_i^3 = -1 \mod 2 \tag{35}$$

and $1 = -1 \mod 2$. We have found that all exotics get a high mass and therefore decouple from the low energy particle spectrum.

The situation of the generation of neutrino masses remains the same also after including the new R symmetries. Thus, neutrino masses can be generated with the see-saw mechanism for many right-handed neutrinos [34, 35].

Our results for proton decay do not differ substantially from the ones described in [4]. We find that $qqql$ operators as well as couplings with massive exotic triplets like $q_1 l_1 \bar{\delta}_4$ and $q_1 q_1 \delta_4$ are allowed. After integrating out the exotic triplets the trilinear operators might be a further source of proton decay. A full clarification of these questions needs a detailed examination of the vacuum configurations of benchmark model 1 and is beyond the scope of this paper.

5 Conclusions

We have shown that with the recently rederived R symmetries the phenomenological attractive models of the mini-landscape remain viable. All appealing features survive and can be understood also from the viewpoint of Hilbert basis monomials. We have shown how symmetries lead to Diophantine equations and how one can find all solutions. It was possible to use this approach for a model in the mini-landscape and its large number of R and non-R symmetries. With this method we have been able to determine the coupling structure to all orders in singlet fields. As an example we have shown that this approach can also be useful for flavor model building and not only for string model building. The symmetries of a given model directly constrain the superpotential and its Hilbert basis building blocks. As has been shown, this can be useful in understanding the F-flatness conditions to find supersymmetry preserving vacua. More work in this direction seems to be interesting. Especially to find point-like minima where all fields and thus all moduli are stabilized. On the other hand the extension of the Hilbert basis approach to non-Abelian discrete symmetries for flavor model building seems to be desirable. An application of our approach to heterotic models based on different orbifold geometries like $\mathbb{Z}_2 \times \mathbb{Z}_2$ is another potential extension.

Acknowledgments

We want to thank Damián Mayorga Peña for discussions. This work was supported by the SFB-Transregio TR33 "The Dark Universe" (Deutsche Forschungsgemeinschaft).

A Details of model 1

Model 1 is based on the gauge embedding

$$\begin{aligned} V &= (\ \tfrac{1}{3},-\tfrac{1}{2},-\tfrac{1}{2},\ \ 0,\ \ 0,\ \ 0,\ \ 0,\ \ 0)(\ \tfrac{1}{2},-\tfrac{1}{6},-\tfrac{1}{2},-\tfrac{1}{2},-\tfrac{1}{2},-\tfrac{1}{2},-\tfrac{1}{2},\ \ \tfrac{1}{2})\,, \\ W_3 = W_4 &= (-\tfrac{1}{2},-\tfrac{1}{2},\ \ \tfrac{1}{6},\ \ \tfrac{1}{6},\ \ \tfrac{1}{6},\ \ \tfrac{1}{6},\ \ \tfrac{1}{6},\ \ \tfrac{1}{6})(\ \tfrac{1}{3},\ \ 0,\ \ 0,\ \ \tfrac{2}{3},\ \ 0,\ \ \tfrac{5}{3},-2,\ \ 0)\,, \\ W_5 &= (\ \ 0,-\tfrac{1}{2},-\tfrac{1}{2},-\tfrac{1}{2},\ \ \tfrac{1}{2},\ \ 0,\ \ 0,\ \ 0)(\ \ 4,-3,-\tfrac{7}{2},-4,-3,-\tfrac{7}{2},-\tfrac{9}{2},\ \ \tfrac{7}{2})\,. \end{aligned}$$

Here we list the complete spectrum of massless string states including their R charges. Those R charges that differ from the "old" ones are marked by red color.

label	k	n_3	n_2	n_2'	q_γ	R^1	R^2	R^3	representation	q_Y	q^1	q^2	q^3	q^4	q^5	q^6	q^7	q^8	q_{anom}	$q_{\text{B}-\text{L}}$
$\bar{n}_3$	0	0	0	0	0	-1	0	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{5}{2}$	0	0	0	0	$-\frac{1}{3}$	-1
$\bar{e}_3$	0	0	0	0	0	-1	0	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	-1	$-\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0	$\frac{2}{3}$	-1
$\bar{u}_3$	0	0	0	0	0	-1	0	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{2}{3}$	$-\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0	$\frac{2}{3}$	$\frac{1}{3}$
$\bar{f}_1$	0	0	0	0	0	-1	0	0	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	0	0	0	0	0	$-\frac{1}{2}$	1	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{5}{3}$	1
f_1	0	0	0	0	0	-1	0	0	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	0	0	0	0	0	$-\frac{1}{2}$	-1	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{2}{3}$	-1
ϕ_1	0	0	0	0	0	0	0	-1	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	-1	1	0	0	0	0	2	0
$\bar{\phi}_1$	0	0	0	0	0	0	0	-1	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	1	-1	0	0	0	0	-2	0
s_2^0	0	0	0	0	0	0	-1	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	0	0	0	0	-1	0	-1	0	$-\frac{5}{3}$	0
s_1^0	0	0	0	0	0	0	-1	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	0	0	0	0	-1	0	1	0	$\frac{7}{3}$	0
q_3	0	0	0	0	0	0	-1	0	$(\mathbf{3},\mathbf{2},\mathbf{1},\mathbf{1})$	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0	$\frac{4}{3}$	$-\frac{1}{3}$
n_{12}	2	0	0	0	0	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{5}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{1}{9}$	1
$\bar{f}_4$	2	0	0	0	0	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{8}{9}$	0
δ_6	2	0	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$-\frac{1}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$-\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{2}{3}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{1}{9}$	$\frac{2}{3}$
$\bar{n}_9$	2	0	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{2}{3}$	$\frac{2}{3}$	0	$-\frac{2}{3}$	$-\frac{1}{9}$	-1
$\bar{\eta}_3$	2	0	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$-\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	$-\frac{2}{3}$	$-\frac{1}{9}$	-1
$\bar{d}_3$	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{4}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{8}{9}$	$\frac{1}{3}$
s_{31}^0	2	0	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{5}{3}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{8}{9}$	0
δ_4	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$-\frac{1}{3}$	0	0	-1	$\frac{2}{3}$	0	0	0	$-\frac{7}{9}$	$\frac{2}{3}$
h_8	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	-1	0	0	$\frac{2}{9}$	0
$\bar{\delta}_4$	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	0	1	$\frac{2}{3}$	0	0	0	$-\frac{1}{9}$	$-\frac{2}{3}$
h_7	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	1	0	0	$\frac{8}{9}$	0
s_{25}^0	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	0	-1	0	$-\frac{13}{9}$	0
s_{24}^0	2	0	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	0	1	0	$\frac{23}{9}$	0
s_{30}^0	2	0	0	0	$\frac{1}{2}$	$\frac{4}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$\frac{2}{3}$	0	0	0	$\frac{2}{9}$	0
s_{26}^0	2	0	0	0	0	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$\frac{2}{3}$	0	0	0	$\frac{2}{9}$	0
$\bar{\delta}_6$	2	2	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$\frac{2}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	$-\frac{1}{3}$	$-\frac{2}{3}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{9}$	$-\frac{2}{3}$
$\bar{n}_{11}$	2	2	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$\frac{4}{3}$	0	$-\frac{1}{3}$	$\frac{5}{9}$	-1
$\bar{n}_{10}$	2	2	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$\frac{1}{2}$	$-\frac{5}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{14}{9}$	-1
$\bar{\eta}_4$	2	2	0	0	0	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{2}{3}$	$\frac{1}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{9}$	-1

label	k	n_3	n_2	n_2'	q_γ	R^1	R^2	R^3	representation	q_Y	q^1	q^2	q^3	q^4	q^5	q^6	q^7	q^8	q_{anom}	$q_{\mathrm{B-L}}$
$\bar{f}_6$	2	2	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$-\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{7}{9}$	0
s_{32}^0	2	2	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$-\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{5}{3}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{2}{9}$	0
$\bar{l}_1$	2	2	0	0	$\frac{1}{2}$	$\frac{7}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{4}{9}$	-1
$\bar{n}_{16}$	2	2	0	0	$\frac{1}{2}$	$\frac{4}{3}$	$-\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{9}$	-1
$\bar{n}_{12}$	2	2	0	0	0	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{9}$	-1
n_{13}	2	0	0	0	1	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{5}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{1}{9}$	1
$\bar{f}_5$	2	0	0	0	1	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{8}{9}$	0
$\bar{d}_4$	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{4}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{8}{9}$	$\frac{1}{3}$
δ_5	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$-\frac{1}{3}$	0	0	-1	$\frac{2}{3}$	0	0	0	$-\frac{7}{9}$	$\frac{2}{3}$
h_{10}	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	-1	0	0	$\frac{2}{9}$	0
$\bar{\delta}_5$	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	0	1	$\frac{2}{3}$	0	0	0	$-\frac{1}{9}$	$-\frac{2}{3}$
h_9	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	1	0	0	$\frac{8}{9}$	0
s_{28}^0	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	0	-1	0	$-\frac{13}{9}$	0
s_{27}^0	2	0	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$-\frac{1}{3}$	0	1	0	$\frac{23}{9}$	0
s_{29}^0	2	0	0	0	1	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	0	0	0	$\frac{2}{3}$	0	0	0	$\frac{2}{9}$	0
$\bar{n}_{14}$	2	2	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$\frac{4}{3}$	0	$-\frac{1}{3}$	$\frac{5}{9}$	-1
$\bar{n}_{13}$	2	2	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$\frac{1}{2}$	$-\frac{5}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{14}{9}$	-1
$\bar{\eta}_5$	2	2	0	0	1	$-\frac{2}{3}$	$\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{2}{3}$	$\frac{1}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{9}$	-1
$\bar{n}_{15}$	2	2	0	0	1	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{9}$	-1
s_{14}^+	3	0	1	1	$\frac{1}{3}$	$-\frac{5}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	$-\frac{1}{2}$	1	$-\frac{1}{2}$	0	$-\frac{5}{6}$	0
s_{14}^-	3	0	1	1	$\frac{1}{3}$	$-\frac{5}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$-\frac{5}{6}$	0
s_{12}^+	3	0	1	1	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
s_{12}^-	3	0	1	1	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	$\frac{1}{2}$	-1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
$\bar{f}_2^+$	3	0	1	1	$-\frac{1}{3}$	$\frac{3}{2}$	0	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	0	0	0	$-\frac{1}{2}$	$-\frac{1}{2}$	-1
$\bar{f}_2^-$	3	0	1	1	$-\frac{1}{3}$	$\frac{3}{2}$	0	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	0	0	0	$\frac{1}{2}$	$\frac{1}{2}$	1
s_{11}^+	3	0	1	0	$\frac{1}{3}$	$-\frac{5}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	$-\frac{1}{2}$	1	$-\frac{1}{2}$	0	$-\frac{5}{6}$	0
s_{11}^-	3	0	1	0	$\frac{1}{3}$	$-\frac{5}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	$-\frac{1}{2}$	-1	$-\frac{1}{2}$	0	$-\frac{5}{6}$	0
s_9^+	3	0	1	0	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
s_9^-	3	0	1	0	0	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	$\frac{1}{2}$	-1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
$\bar{f}_1^+$	3	0	1	0	$-\frac{1}{3}$	$\frac{3}{2}$	0	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	0	0	0	$-\frac{1}{2}$	$-\frac{1}{2}$	-1
$\bar{f}_1^-$	3	0	1	0	$-\frac{1}{3}$	$\frac{3}{2}$	0	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	0	0	0	$\frac{1}{2}$	$\frac{1}{2}$	1
h_6	3	0	0	1	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	1	0	0	$-\frac{1}{2}$	0
h_5	3	0	0	1	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	-1	0	0	$\frac{1}{2}$	0
χ_4	3	0	0	1	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0	1	$-\frac{1}{2}$	2
χ_3	3	0	0	1	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	0	-1	$\frac{1}{2}$	-2
h_4	3	0	0	0	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	1	0	0	$-\frac{1}{2}$	0
h_3	3	0	0	0	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	-1	0	0	$\frac{1}{2}$	0
χ_2	3	0	0	0	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	0	$-\frac{1}{2}$	$\frac{1}{2}$	0	0	0	0	1	$-\frac{1}{2}$	2
χ_1	3	0	0	0	$-\frac{1}{3}$	$\frac{3}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	0	-1	$\frac{1}{2}$	-2
s_{13}^+	3	0	1	1	1	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
s_{13}^-	3	0	1	1	1	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	$\frac{1}{2}$	-1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
s_{10}^+	3	0	1	0	1	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	0	0	0	-1	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
s_{10}^-	3	0	1	0	1	$-\frac{1}{2}$	0	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	0	0	0	1	$\frac{1}{2}$	-1	$\frac{1}{2}$	0	$\frac{5}{6}$	0
s_{22}^0	4	2	0	0	0	$-\frac{1}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{5}{3}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{9}$	0
f_5	4	2	0	0	0	$-\frac{1}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{7}{9}$	0

label	k	n_3	n_2	n_2'	q_γ	R^1	R^2	R^3	representation	q_Y	q^1	q^2	q^3	q^4	q^5	q^6	q^7	q^8	q_{anom}	$q_{\mathrm{B-L}}$
l_3	4	2	0	0	0	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{4}{9}$	1
n_{10}	4	2	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$-\frac{1}{2}$	$\frac{5}{6}$	$-\frac{5}{6}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{14}{9}$	1
n_9	4	2	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{3}$	$-\frac{4}{3}$	0	$\frac{1}{3}$	$-\frac{5}{9}$	1
η_5	4	2	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$-\frac{2}{3}$	$-\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{1}{9}$	1
δ_2	4	2	0	0	0	$-\frac{1}{3}$	$\frac{4}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{1}{9}$	$\frac{2}{3}$
n_7	4	2	0	0	0	$\frac{2}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{1}{9}$	1
n_{11}	4	2	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{1}{9}$	1
h_1	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$-\frac{2}{3}$	0	0	0	$\frac{1}{3}$	1	0	0	$-\frac{2}{9}$	0
$\bar{\delta}_1$	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{3}$	0	0	1	$-\frac{2}{3}$	0	0	0	$\frac{7}{9}$	$-\frac{2}{3}$
h_2	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$-\frac{2}{3}$	0	0	0	$\frac{1}{3}$	-1	0	0	$-\frac{8}{9}$	0
δ_1	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$\frac{1}{3}$	0	0	-1	$-\frac{2}{3}$	0	0	0	$\frac{1}{9}$	$\frac{2}{3}$
s_{18}^0	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	0	0	$\frac{1}{3}$	0	-1	0	$-\frac{23}{9}$	0
s_{17}^0	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	0	0	$\frac{1}{3}$	0	1	0	$\frac{13}{9}$	0
s_{15}^0	4	0	0	0	0	$\frac{2}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	0	0	$-\frac{2}{3}$	0	0	0	$-\frac{2}{9}$	0
s_{19}^0	4	0	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	0	0	$-\frac{2}{3}$	0	0	0	$-\frac{2}{9}$	0
s_{20}^0	4	1	0	0	0	$-\frac{1}{3}$	$\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{5}{3}$	$\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{8}{9}$	0
d_1	4	1	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$\frac{1}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$-\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{6}$	$\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{8}{9}$	$-\frac{1}{3}$
η_3	4	1	0	0	0	$-\frac{1}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{1}{3}$	$\frac{1}{3}$	0	$\frac{2}{3}$	$\frac{1}{9}$	1
f_4	4	1	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{1}{6}$	$\frac{1}{3}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{8}{9}$	0
$\bar{n}_8$	4	1	0	0	$\frac{1}{2}$	$\frac{8}{3}$	$\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{5}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{1}{9}$	-1
n_5	4	1	0	0	0	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{2}{3}$	$-\frac{2}{3}$	0	$\frac{2}{3}$	$\frac{1}{9}$	1
$\bar{\delta}_2$	4	1	0	0	0	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{3}$	0	$-\frac{1}{3}$	$-\frac{2}{3}$	$\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{1}{9}$	$-\frac{2}{3}$
s_{23}^0	4	2	0	0	1	$-\frac{1}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{5}{3}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$-\frac{2}{9}$	0
f_6	4	2	0	0	1	$-\frac{1}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{7}{9}$	0
l_4	4	2	0	0	1	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{4}{9}$	1
δ_3	4	2	0	0	1	$-\frac{1}{3}$	$\frac{4}{3}$	0	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{3}$	$\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{1}{9}$	$\frac{2}{3}$
n_8	4	2	0	0	1	$\frac{2}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{1}{9}$	1
s_{16}^0	4	0	0	0	1	$\frac{2}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	0	0	$-\frac{2}{3}$	0	0	0	$-\frac{2}{9}$	0
s_{21}^0	4	1	0	0	1	$-\frac{1}{3}$	$\frac{4}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{5}{3}$	$\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{8}{9}$	0
η_4	4	1	0	0	1	$-\frac{1}{3}$	$\frac{1}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{1}{3}$	$\frac{1}{3}$	0	$\frac{2}{3}$	$\frac{1}{9}$	1
n_6	4	1	0	0	1	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$-\frac{2}{3}$	$-\frac{2}{3}$	0	$\frac{2}{3}$	$\frac{1}{9}$	1
$\bar{\delta}_3$	4	1	0	0	1	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{3}$	0	$-\frac{1}{3}$	$-\frac{2}{3}$	$\frac{1}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{1}{9}$	$-\frac{2}{3}$
s_4^+	5	1	1	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$-\frac{1}{3}$	0	$-\frac{2}{3}$	$\frac{2}{3}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{37}{18}$	0
ν_2	5	1	1	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{6}$	$-\frac{1}{3}$	0	$\frac{1}{3}$	$-\frac{1}{3}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{18}$	$\frac{2}{3}$
s_3^-	5	1	1	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{2}{3}$	$-\frac{17}{18}$	-1
m_6	5	1	1	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{19}{18}$	1
s_3^+	5	1	1	1	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{19}{18}$	0
s_4^-	5	1	1	1	0	$\frac{5}{6}$	$\frac{2}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{25}{18}$	1
s_2^+	5	1	1	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$-\frac{1}{3}$	0	$-\frac{2}{3}$	$\frac{2}{3}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{37}{18}$	0
nu_1	5	1	1	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{3},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{6}$	$-\frac{1}{3}$	0	$\frac{1}{3}$	$-\frac{1}{3}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{18}$	$\frac{2}{3}$
s_1^-	5	1	1	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{2}{3}$	$-\frac{17}{18}$	-1
m_5	5	1	1	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$\frac{1}{2}$	$-\frac{1}{6}$	$-\frac{5}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{19}{18}$	1
s_1^+	5	1	1	0	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{19}{18}$	0
s_2^-	5	1	1	0	0	$\frac{5}{6}$	$\frac{2}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{25}{18}$	1
η_2	5	1	0	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	0	$-\frac{2}{3}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{19}{18}$	1

label	k	n_3	n_2	n_2'	q_γ	R^1	R^2	R^3	representation	q_Y	q^1	q^2	q^3	q^4	q^5	q^6	q^7	q^8	q_{anom}	$q_{\mathrm{B-L}}$
$\bar{n}_5$	5	1	0	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{2}{3}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$-\frac{2}{3}$	$\frac{19}{18}$	-1
n_2	5	1	0	1	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{2}{3}$	$-\frac{5}{6}$	$\frac{2}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{19}{18}$	1
η_1	5	1	0	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	0	$-\frac{2}{3}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{19}{18}$	1
n_1	5	1	0	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{2}{3}$	$-\frac{5}{6}$	$-\frac{1}{3}$	$\frac{2}{3}$	0	$-\frac{2}{3}$	$\frac{19}{18}$	-1
$\bar{n}_4$	5	1	0	0	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{2}{3}$	$-\frac{5}{6}$	$\frac{2}{3}$	$\frac{2}{3}$	0	$\frac{1}{3}$	$\frac{19}{18}$	1
y_2	5	0	1	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{2})$	0	$-\frac{1}{3}$	0	0	0	$\frac{1}{6}$	0	$\frac{1}{2}$	0	$\frac{13}{18}$	0
m_3	5	0	1	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	0	0	0	$\frac{1}{6}$	1	$-\frac{1}{2}$	0	$-\frac{17}{18}$	0
m_4	5	0	1	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	0	0	0	$\frac{1}{6}$	-1	$-\frac{1}{2}$	0	$-\frac{29}{18}$	0
y_1	5	0	1	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{2})$	0	$-\frac{1}{3}$	0	0	0	$\frac{1}{6}$	0	$\frac{1}{2}$	0	$\frac{13}{18}$	0
m_1	5	0	1	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	0	0	0	$\frac{1}{6}$	1	$-\frac{1}{2}$	0	$-\frac{17}{18}$	0
m_2	5	0	1	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	0	0	0	$\frac{1}{6}$	-1	$-\frac{1}{2}$	0	$-\frac{29}{18}$	0
$\bar{d}_1$	5	0	0	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{6}$	0	0	$-\frac{3}{2}$	$-\frac{1}{3}$	0	0	0	$-\frac{5}{18}$	$\frac{1}{3}$
$\bar{e}_1$	5	0	0	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	-1	$\frac{1}{6}$	0	0	$\frac{1}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{7}{18}$	-1
$\bar{u}_1$	5	0	0	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{2}{3}$	$\frac{1}{6}$	0	0	$\frac{1}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{7}{18}$	$\frac{1}{3}$
l_1	5	0	0	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{1}{6}$	0	0	$-\frac{3}{2}$	$-\frac{1}{3}$	0	0	0	$-\frac{5}{18}$	1
q_1	5	0	0	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{3},\mathbf{2},\mathbf{1},\mathbf{1})$	$-\frac{1}{6}$	$\frac{1}{6}$	0	0	$\frac{1}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{7}{18}$	$-\frac{1}{3}$
$\bar{n}_1$	5	0	0	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	0	$\frac{5}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{19}{18}$	-1
s_{12}^0	5	0	0	1	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$-\frac{5}{18}$	0
s_{11}^0	5	0	0	1	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	$\frac{1}{2}$	$-\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$\frac{25}{18}$	0
s_{14}^0	5	0	0	1	0	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$\frac{13}{18}$	0
s_{13}^0	5	0	0	1	0	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$-\frac{17}{18}$	0
s_{10}^0	5	0	0	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$\frac{13}{18}$	0
s_9^0	5	0	0	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$-\frac{17}{18}$	0
$\bar{d}_2$	5	0	0	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{3}$	$\frac{1}{6}$	0	0	$-\frac{3}{2}$	$-\frac{1}{3}$	0	0	0	$-\frac{5}{18}$	$\frac{1}{3}$
$\bar{e}_2$	5	0	0	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	-1	$\frac{1}{6}$	0	0	$\frac{1}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{7}{18}$	-1
$\bar{u}_2$	5	0	0	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{2}{3}$	$\frac{1}{6}$	0	0	$\frac{1}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{7}{18}$	$\frac{1}{3}$
l_2	5	0	0	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{1}{6}$	0	0	$-\frac{3}{2}$	$-\frac{1}{3}$	0	0	0	$-\frac{5}{18}$	1
q_2	5	0	0	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{3},\mathbf{2},\mathbf{1},\mathbf{1})$	$-\frac{1}{6}$	$\frac{1}{6}$	0	0	$\frac{1}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{7}{18}$	$-\frac{1}{3}$
$\bar{n}_2$	5	0	0	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	0	$\frac{5}{2}$	$-\frac{1}{3}$	0	0	0	$\frac{19}{18}$	-1
s_6^0	5	0	0	0	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	$-\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$-\frac{5}{18}$	0
s_5^0	5	0	0	0	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{2}{3}$	$\frac{1}{2}$	$-\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$\frac{25}{18}$	0
s_8^0	5	0	0	0	0	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$\frac{13}{18}$	0
s_7^0	5	0	0	0	0	$\frac{11}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$-\frac{17}{18}$	0
s_4^0	5	0	0	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$-\frac{1}{2}$	$-\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$\frac{13}{18}$	0
s_3^0	5	0	0	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$-\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{3}$	0	0	0	$-\frac{17}{18}$	0
$\bar{\nu}_2$	5	1	1	1	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{6}$	$-\frac{1}{3}$	0	$-\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{25}{18}$	$-\frac{2}{3}$
s_8^-	5	1	1	1	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$-\frac{1}{3}$	0	$\frac{2}{3}$	$-\frac{2}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$-\frac{11}{18}$	0
s_7^+	5	1	1	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$-\frac{1}{2}$	$\frac{2}{3}$	$-\frac{17}{18}$	1
m_8	5	1	1	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{19}{18}$	-1
s_7^-	5	1	1	1	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{2}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{31}{18}$	0
s_8^+	5	1	1	1	0	$\frac{5}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{13}{18}$	-1
$\bar{\nu}_1$	5	1	1	0	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\bar{\mathbf{3}},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{6}$	$-\frac{1}{3}$	0	$-\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{25}{18}$	$-\frac{2}{3}$
s_6^-	5	1	1	0	0	$-\frac{1}{6}$	$-\frac{4}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$-\frac{1}{3}$	0	$\frac{2}{3}$	$-\frac{2}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$-\frac{11}{18}$	0
s_5^+	5	1	1	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$-\frac{1}{2}$	$\frac{2}{3}$	$-\frac{17}{18}$	1
m_7	5	1	1	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{2},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{19}{18}$	-1

label	k	n_3	n_2	n_2'	q_γ	R^1	R^2	R^3	representation	q_Y	q^1	q^2	q^3	q^4	q^5	q^6	q^7	q^8	q_{anom}	$q_{\mathrm{B-L}}$
s_5^-	5	1	1	0	0	$-\frac{1}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$\frac{1}{2}$	$\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{2}{3}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{31}{18}$	0
s_6^+	5	1	1	0	0	$\frac{5}{6}$	$-\frac{4}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$-\frac{1}{3}$	$\frac{13}{18}$	-1
f_3	5	1	0	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{1}{18}$	0
$\bar{f}_3$	5	1	0	1	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{37}{18}$	0
$\bar{\eta}_2$	5	1	0	1	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$-\frac{1}{3}$	$\frac{1}{3}$	0	$-\frac{1}{3}$	$\frac{19}{18}$	-1
n_4	5	1	0	1	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{2}{3}$	$\frac{19}{18}$	1
$\bar{n}_7$	5	1	0	1	0	$\frac{5}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$\frac{2}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{7}{18}$	-1
f_2	5	1	0	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{4},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$-\frac{1}{2}$	$\frac{1}{6}$	$\frac{1}{18}$	0
$\bar{f}_2$	5	1	0	0	0	$-\frac{1}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\bar{\mathbf{4}},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{2}$	$\frac{1}{6}$	$\frac{37}{18}$	0
$\bar{\eta}_1$	5	1	0	0	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{2})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$-\frac{1}{3}$	$\frac{1}{3}$	0	$-\frac{1}{3}$	$\frac{19}{18}$	-1
n_3	5	1	0	0	0	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$-\frac{1}{3}$	$-\frac{2}{3}$	0	$\frac{2}{3}$	$\frac{19}{18}$	1
$\bar{n}_6$	5	1	0	0	0	$\frac{5}{6}$	$\frac{2}{3}$	$-\frac{1}{2}$	$(\mathbf{1},\mathbf{1},\mathbf{1},\mathbf{1})$	0	$\frac{1}{6}$	0	$-\frac{1}{3}$	$\frac{5}{6}$	$\frac{2}{3}$	$-\frac{2}{3}$	0	$-\frac{1}{3}$	$\frac{7}{18}$	-1

References

[1] H. P. Nilles, S. Ramos-Sanchez, M. Ratz, and P. K. Vaudrevange, "A note on discrete R symmetries in $\mathbb{Z}_6$-II orbifolds with Wilson lines," *Phys.Lett.* **B726** (2013) 876–881, arXiv:1308.3435 [hep-th].

[2] N. G. Cabo Bizet, T. Kobayashi, D. K. Mayorga Pena, S. L. Parameswaran, M. Schmitz, *et al.*, "Discrete R-symmetries and Anomaly Universality in Heterotic Orbifolds," *JHEP* **1402** (2014) 098, arXiv:1308.5669 [hep-th].

[3] O. Lebedev, H. P. Nilles, S. Raby, S. Ramos-Sanchez, M. Ratz, *et al.*, "A Mini-landscape of exact MSSM spectra in heterotic orbifolds," *Phys.Lett.* **B645** (2007) 88–94, arXiv:hep-th/0611095 [hep-th].

[4] O. Lebedev, H. P. Nilles, S. Raby, S. Ramos-Sanchez, M. Ratz, *et al.*, "The Heterotic Road to the MSSM with R parity," *Phys.Rev.* **D77** (2008) 046013, arXiv:0708.2691 [hep-th].

[5] O. Lebedev, H. P. Nilles, S. Ramos-Sanchez, M. Ratz, and P. K. Vaudrevange, "Heterotic mini-landscape. (II). Completing the search for MSSM vacua in a Z(6) orbifold," *Phys.Lett.* **B668** (2008) 331–335, arXiv:0807.4384 [hep-th].

[6] R. Kappl, M. Ratz, and C. Staudt, "The Hilbert basis method for D-flat directions and the superpotential," *JHEP* **1110** (2011) 027, arXiv:1108.2154 [hep-th].

[7] R. Kappl, B. Petersen, S. Raby, M. Ratz, R. Schieren, *et al.*, "String-Derived MSSM Vacua with Residual R Symmetries," *Nucl.Phys.* **B847** (2011) 325–349, arXiv:1012.4574 [hep-th].

[8] S. Antusch, I. de Medeiros Varzielas, V. Maurer, C. Sluka, and M. Spinrath, "Towards predictive flavour models in SUSY SU(5) GUTs with doublet-triplet splitting," arXiv:1405.6962 [hep-ph].

[9] W. Bruns, B. Ichim, T. Römer, and C. Söger, "Normaliz. Algorithms for rational cones and affine monoids." Available from http://www.math.uos.de/normaliz.

[10] 4ti2 team, "4ti2—a software package for algebraic, geometric and combinatorial problems on linear spaces." Available at www.4ti2.de.

[11] W. Bruns, B. Ichim, and C. Söger, "The power of pyramid decomposition in Normaliz," *ArXiv e-prints* (June, 2012) , arXiv:1206.1916 [math.CO].

[12] M.-C. Chen, J. Huang, J.-M. O'Bryan, A. M. Wijangco, and F. Yu, "Compatibility of θ_{13} and the Type I Seesaw Model with A_4 Symmetry," *JHEP* **1302** (2013) 021, arXiv:1210.6982 [hep-ph].

[13] W. Buchmuller, K. Hamaguchi, O. Lebedev, and M. Ratz, "Supersymmetric Standard Model from the Heterotic String (II)," *Nucl.Phys.* **B785** (2007) 149–209, arXiv:hep-th/0606187 [hep-th].

[14] W. Bruns and B. Ichim, "Normaliz: Algorithms for affine monoids and rational cones," *Journal of Algebra* **324** no. 5, (2010) 1098 – 1113. Computational Algebra.

[15] R. Kappl, H. P. Nilles, S. Ramos-Sanchez, M. Ratz, K. Schmidt-Hoberg, *et al.*, "Large hierarchies from approximate R symmetries," *Phys.Rev.Lett.* **102** (2009) 121602, arXiv:0812.2120 [hep-th].

[16] F. Brummer, R. Kappl, M. Ratz, and K. Schmidt-Hoberg, "Approximate R-symmetries and the mu term," *JHEP* **1004** (2010) 006, arXiv:1003.0084 [hep-th].

[17] P. Ko, T. Kobayashi, J.-h. Park, and S. Raby, "String-derived D(4) flavor symmetry and phenomenological implications," *Phys.Rev.* **D76** (2007) 035005, arXiv:0704.2807 [hep-ph].

[18] T. Kobayashi, H. P. Nilles, F. Ploger, S. Raby, and M. Ratz, "Stringy origin of non-Abelian discrete flavor symmetries," *Nucl.Phys.* **B768** (2007) 135–156, arXiv:hep-ph/0611020 [hep-ph].

[19] N. G. Cabo Bizet, T. Kobayashi, D. K. Mayorga Pena, S. L. Parameswaran, M. Schmitz, *et al.*, "R-charge Conservation and More in Factorizable and Non-Factorizable Orbifolds," *JHEP* **1305** (2013) 076, arXiv:1301.2322 [hep-th].

[20] H. P. Nilles, S. Ramos-Sanchez, P. K. Vaudrevange, and A. Wingerter, "The Orbifolder: A Tool to study the Low Energy Effective Theory of Heterotic Orbifolds," *Comput.Phys.Commun.* **183** (2012) 1363–1380, arXiv:1110.5229 [hep-th].

[21] M. Maniatis, A. von Manteuffel, and O. Nachtmann, "Determining the global minimum of Higgs potentials via Groebner bases: Applied to the NMSSM," *Eur.Phys.J.* **C49** (2007) 1067–1076, arXiv:hep-ph/0608314 [hep-ph].

[22] J. Gray, Y.-H. He, and A. Lukas, "Algorithmic Algebraic Geometry and Flux Vacua," *JHEP* **0609** (2006) 031, arXiv:hep-th/0606122 [hep-th].

[23] J. Gray, Y.-H. He, A. Ilderton, and A. Lukas, "A New Method for Finding Vacua in String Phenomenology," *JHEP* **0707** (2007) 023, arXiv:hep-th/0703249 [HEP-TH].

[24] W. Decker, G.-M. Greuel, G. Pfister, and H. Schönemann, "SINGULAR 3-1-6 — A computer algebra system for polynomial computations." http://www.singular.uni-kl.de, 2012.

[25] H. M. Lee, S. Raby, M. Ratz, G. G. Ross, R. Schieren, *et al.*, "A unique Z_4^R symmetry for the MSSM," *Phys.Lett.* **B694** (2011) 491–495, arXiv:1009.0905 [hep-ph].

[26] H. K. Dreiner, T. Opferkuch, and C. Luhn, "Froggatt-Nielsen models with a residual $\mathbb{Z}_4^R$ symmetry," *Phys.Rev.* **D88** no. 11, (2013) 115005, arXiv:1308.0332 [hep-ph].

[27] M.-C. Chen, M. Ratz, and V. Takhistov, "R parity violation from discrete R symmetries," arXiv:1410.3474 [hep-ph].

[28] J. Camargo-Molina, B. O'Leary, W. Porod, and F. Staub, "**Vevacious**: A Tool For Finding The Global Minima Of One-Loop Effective Potentials With Many Scalars," *Eur.Phys.J.* **C73** (2013) 2588, arXiv:1307.1477 [hep-ph].

[29] M. Cicoli, D. Klevers, S. Krippendorf, C. Mayrhofer, F. Quevedo, *et al.*, "Explicit de Sitter Flux Vacua for Global String Models with Chiral Matter," *JHEP* **1405** (2014) 001, arXiv:1312.0014 [hep-th].

[30] F. Buccella, J. Derendinger, S. Ferrara, and C. A. Savoy, "Patterns of Symmetry Breaking in Supersymmetric Gauge Theories," *Phys.Lett.* **B115** (1982) 375.

[31] G. Cleaver, M. Cvetic, J. R. Espinosa, L. L. Everett, and P. Langacker, "Classification of flat directions in perturbative heterotic superstring vacua with anomalous U(1)," *Nucl.Phys.* **B525** (1998) 3–26, arXiv:hep-th/9711178 [hep-th].

[32] P. Hosteins, R. Kappl, M. Ratz, and K. Schmidt-Hoberg, "Gauge-top unification," *JHEP* **0907** (2009) 029, arXiv:0905.3323 [hep-ph].

[33] R. Kappl, "Quark mass hierarchies in heterotic orbifold GUTs," *JHEP* **1104** (2011) 019, arXiv:1012.4368 [hep-ph].

[34] W. Buchmuller, K. Hamaguchi, O. Lebedev, S. Ramos-Sanchez, and M. Ratz, "Seesaw neutrinos from the heterotic string," *Phys.Rev.Lett.* **99** (2007) 021601, arXiv:hep-ph/0703078 [HEP-PH].

[35] J. Heeck, "Seesaw parametrization for n right-handed neutrinos," *Phys.Rev.* **D86** (2012) 093023, arXiv:1207.5521 [hep-ph].